York Foundation Trust R&D Unit Standard Operating Procedure R&D/S21



Study Close-Out: Clinical Trials of Investigational Medicinal Products (CTIMPs) and other research studies

IT IS THE RESPONSIBILITY OF <u>ALL</u> USERS OF THIS SOP TO ENSURE THAT THE CORRECT VERSION IS BEING USED

All staff should regularly check the R&D Unit's website and/or Q-Pulse for information relating to the implementation of new or revised versions. Staff must ensure that they are adequately trained in the new procedure and must make sure that all copies of superseded versions are promptly withdrawn from use unless notified otherwise by the SOP Controller.

The definitive versions of all R&D Unit SOPs appear online. If you are reading this in printed form check that the version number and date below is the most recent one as shown on the R&D Unit website: www.northyorksresearch.nhs.uk/sops.html and/or Q-Pulse

SOP Reference: R&D/S21 Version Number: 3.0

Signature:

Author: Tracey Dorey Implementation date of current version: 21st August 2017

Approved by: Name/Position: Lydia Harris, Head of R&D

Date: 18th July 2017

Name/Position: Sarah Sheath, SOP Controller

Signed copy held by R&D Unit

Signature: Signed copy held by R&D Unit

Date: 18th July 2017

This SOP will normally be reviewed every 3 years unless changes to the legislation require otherwise

© York Teaching Hospital NHS Foundation Trust 2017 All Rights Reserved

Version History Log

This area should detail the version history for this document. It should detail the key elements of the changes to the versions.

| Version | Date Implemented | Details of significant changes |
|-------------|-------------------------------|---|
| 1.0 | 1 st November 2010 | |
| 2.0 | 17 th October 2013 | Change of SOP Controller and author. Removal of references to the North and East Yorkshire R&D Alliance |
| | | Made applicable to other research in addition to |
| | | CTIMP studies |
| 3.0 | 21 st August 2017 | Change of author, routine review, minor change. |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | (,) |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| |) ` | |
| | | |
| 1/19 | | |
| <i>\)</i> . | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |

Version 3.0 Contents

Contents

Page No

| 1 | Introduction, Background and Purpose | | |
|---|---|------------------|--|
| 2 | 2 Who Should Use This SOP | | |
| 3 | When this SOP Should be Used | | |
| 4 | Procedure(s) | 2 | |
| | 4.1 Chief Investigator (CI) Responsibilities – Trust Sponsored Studies 4.2 Pharmacy 4.3 Clinical Samples 4.4 Close-Out Monitoring Visit 4.5 R&D Unit Responsibilities | 2 3 3 3 | |
| 5 | Related SOPs and Documents | 3 | |

Version 3.0 Contents

1 Introduction, Background and Purpose

Trial Close-out is the act of ensuring that all clinical trial related activities have been appropriately reconciled.

Close-out is integral to the quality of a trial and is designed to ensure that all necessary documents are in place should the trial data need to be queried or inspected in the future.

This SOP establishes a procedure for closing down clinical trials of investigational medicinal products (CTIMPs) in the Trust to support compliance with the UK Clinical Trial Regulations. A similar approach should be adopted for non-CTIMP studies to ensure consistent practice.

2 Who Should Use This SOP

This SOP is aimed at:

- Chief Investigators (CIs), trial co-ordinators, and other professional or administrative research staff, in the investigator team or in support departments, working on CTIMP and non-CTIMP studies Sponsored or co-sponsored by the Trust;
- Principal Investigators (PIs) and other professional or administrative research staff, in the investigator team or in support departments, at Sites where multi-site studies sponsored or co-sponsored by the Trust are being run;
- R&D Unit personnel, who manage the sponsorship of CTIMP and non-CTIMP studies on behalf of the Trust;
- Pls, research staff and staff in all involved service departments in the Trust where externally-sponsored studies are hosted;
- Monitors employed by or contracted to the sponsoring or co-sponsoring the Trust.

3 When this SOP Should be Used

This SOP should be followed when closing down:

- a CTIMP or non-CTIMP study that is sponsored or co-sponsored by the Trust;
- a CTIMP hosted by the Trust with an external Sponsor, where an applicable Sponsor SOP has not been adopted for the trial.

This SOP begins when the study terminates. The end of study should be defined in the study protocol but may be for example, when the last patient entered onto the study has had their last study visit. Any changes to the end of study should be notified to the authorities as a Substantial Amendment. Final analysis of the data (following 'lock' of the trial database) and report writing may occur after formal declaration of the end of the project.

Version 2.0 Page 1 of 4

4 Procedure(s)

A site must be closed as soon as is practicable to do so. A site may be deemed "closed" once all study-related activities at a particular site are reconciled and/or complete. This includes ensuring:

- Investigator/institution and sponsor files are reviewed and all essential documentation for a particular site are confirmed in the appropriate files, providing a clear audit trail of study conduct at the site.
- All site data is collected, entered, validated and all data queries resolved where feasible. This includes queries resulting from reconciliation of the clinical and safety database.
- All issues from previous study monitoring procedures are resolved and documented.
- All financial matters are resolved and all site payments are complete as agreed and documented in study contracts/agreements/approvals.
- All unused trial supplies are returned or destroyed according to study and/or sponsor requirements.
- Final drug accountability is complete and return (if returned) or destruction of unused study drug is documented in the site file (if destroyed locally at site).
- Investigator(s) are aware of the study publication policy, as documented in the study protocol and/or study contracts/agreements.
- Investigator(s) are aware of and have implemented relevant ongoing requirements such as site archiving, subsequent audit/inspection procedures and any ongoing reporting requirements.

The Chief Investigator (CI) site should not be closed until all the participating sites have been closed-out.

Consideration must also be given to those trials that can be 'closed' but where the patients are placed into long term follow-up (e.g. Oncology trials).

4.1 Chief Investigator (CI) Responsibilities – Trust Sponsored Studies

- It is the responsibility of the CI, acting on behalf of the Sponsor, to notify the main REC, the MHRA (CTIMPs only) and the Sponsor (via R&D Unit) of the end of the trial for every participating site and CI site using the Declaration of End of Trial Form (see R&D/S06 for details).
- The CI must notify the end of the studyl within 90 days of the trial ending (trial end as defined in the protocol) or within 15 days if terminated early.
- Once notified of the end of study, the CI will contact the Monitor in order to arrange a Close-Out monitoring visit for each site. Support departments (e.g. pharmacy) should also be notified in order that they can prepare for Close-Out.
- Organise Trial Master File (TMF) ensuring all necessary documents are present.
- The CI is must submit an end of study report to the R&D Unit within 10 months of the date of the end of trial. This Report will be submitted to

Version 2.0 Page 2 of 4

the York Teaching Hospital NHS Foundation Trust R&D Group for information. The R&D Unit will subsequently be responsible for submitting the end of study report to the MHRA (CTIMPS only) and Main Ethics Committee to arrive within 12 months of the date of the end of the study (see R&D/S27).

 Instigate archiving procedures in line with R&D/S11 ensuring that files are retrieved from support departments (e.g. pharmacy) so that all essential documents relating to a particular trial are archived together.

4.2 Pharmacy

Pharmacy Close-Out procedures are covered in more detail in Pharm/S56 (Close Down of a Clinical Trial in Pharmacy).

All Investigational Medicinal Product (IMP) must be accounted for, including a check that all IMP has been returned by trial subjects and that all IMP not used for the study is present and unopened. All IMP logs must be checked for accuracy and any discrepancies must be accounted for. Serious breaches in IMP accountability must be brought to the attention of the CI and Sponsor.

Return and destruction of IMPs must be carried out as according to protocol and legal requirements. IMPs must not be destroyed until permission to do so is given by the Sponsor. A Certificate of Destruction should be obtained when the IMP is destroyed, this should be filed in the Trial Pharmacy File.

4.3 Clinical Samples

Storage and destruction of clinical samples must be carried out according to Protocol and Sponsor requirements.

4.4 Close-Out Monitoring Visit

Monitor's Close-Out procedures are covered in more detail in SOP R&D/S08.

4.5 R&D Unit Responsibilities

- 1. Receive study conclusion notification from CI.
- 2. Update end date on EDGE and change study status to completed/closed.
- 3. Notify the CI that the study is closed, once the close-out procedures have been completed.
- 4. Prepare R&D Unit files (and Sponsor files where applicable) for archiving.
- 5. At the time when trial documentation is ready for archive, the Named Archivist will prepare archive boxes in line with SOP R&D/S11.

5 Related SOPs and Documents

R&D/S03 Delegation of Roles and Responsibilities for Trust Sponsored

Research Studies

R&D/S06 Safety Reporting

Version 2.0 Page 3 of 4

| R&D/S08 | Monitoring of Trust sponsored Research Studies |
|----------|--|
| R&D/S09 | Set Up and Management of Research Studies |
| R&D/S11 | Archiving of Essential Documents |
| R&D/S27 | End of Study Reports |
| Pharm/56 | Trial Closedown in Pharmacy |
| R&D/F19 | Close-Out Visit Monitoring Report |
| | ROLLED DOCUMENT WHEN PRINTED |

Version 2.0 Page 4 of 4